

A NEW SPECIES AND A NEW RECORD SPECIES OF THE GENUS *SINGAPORA* MAHMOOD (HEMIPTERA, CICADELLIDAE, TYPHLOCYBINAЕ) FROM CHINA

SONG YueHua^{1, 2}, LIZiZhong^{1*}

1. The Provincial Key Laboratory for Agricultural Pest Management of Mountainous Region, Institute of Entomology, Guizhou University, Guiyang, Guizhou 550025, China

2. Institute of South China Karst, Guizhou Normal University, Guiyang, Guizhou 550001, China

Abstract A new species *S. bannaensis* and a new record species *S. arifi* Ghauri, 1985 are described and illustrated from Yunnan Province, China. A key to males of *Singapora* species is provided.

Key words Hemiptera, Cicadellidae, Typhlocybinae, *Singapora*, new species, China

Introduction

The leafhopper genus *Singapora* Mahmood, 1967 belongs to Typhlocybinae tribe Erythroneurini (type species *S. nigropunctata* Mahmood). It was reviewed by Ghauri (1985) and eleven species are known worldwide so far, including four species from China. A new species and a new record species are described and illustrated from China in this paper. All specimens examined are deposited in the collection of the Institute of Entomology, Guizhou University (IEGU), China.

Singapora Mahmood, 1967

Singapora Mahmood, 1967: 20; Dworakowska, 1970: 759; Dworakowska et Sohj, 1978: 39; Chou et Ma, 1981: 196; Dworakowska, 1983: 118; Ghauri, 1985: 5; Chiang et Knight, 1990: 240.

Erythroneuropsis Ranakrishnan et Menon, 1973: 37.

Type species: *Singapora nigropunctata* Mahmood, 1967.

Body usually yellow, or white, or greenish yellow, or brown. Head narrower than pronotum. Crown anterior margin obtusely rounded, slightly produced medially, subparallel to posterior margin. Vertex with large median apical spot, black or brownish black, which only in female in some species. Eyes black; ocelli present. Face not depressed in profile, 45° or more from horizontal; anteclypeus pale or dark, narrow or inflated slightly. Pronotum usually without conspicuous patterns, but sometimes central region and near hind margin dark. Forewing 1st apical cell largest, with transverse base; 2nd and 3rd apical cells nearly equal in width, 2nd slightly longer than 3rd; 4th apical cell reaching wing apex.

Abdominal apodemes broad, not or exceeding 3rd sternite.

Male genitalia. Anal tube appendage present.

Pygofer without appendages, with a few slender setae scattered on ventral surface of lobe. Subgenital plate gradually curved dorsad, apex rounded, with several macrosetae at about middle of outer margin and with marginal subbasal group of macrosetae. Style with some furrows at apex, midlength of style broad with some gracile microsetae near preapical lobe. Connective Y-shaped, with short central lobe and well developed manubrium. Aedeagal shaft curved dorsad. Aedeagus preatrium with single well developed process, dorsal apodeme distinct.

Distribution: Singapore, North Korea, India (Karnataka, Madhya Pradesh, Haryana, Pilani, Assam), Thailand (Bangkok, Saraburi), USSR (Uzbek); China (Taiwan, Shaanxi, Yunnan, Nanjing, Beijing, Hangzhou, Guangzhou).

Key to males of *Singapora* of the species

1. Pronotum with yellow brownish markings, scutellum without dark spots *S. cyclops* Kuznetsov
Pronotum without yellow brownish markings, or if pronotum with markings, then scutellum also maculate 2
2. Aedeagal shaft with pair of apical processes 3
Aedeagal shaft without apical processes 7
3. Aedeagus with long preatrium process 4
Aedeagus without preatrium process *S. diversa* Ghauri
4. Apical processes of aedeagus short, not extending to 1/2 length of shaft 5
Apical processes of aedeagus long, extending beyond 1/2 length of shaft 6
5. Aedeagal shaft very long and slim, curved obviously, C-shaped
..... *S. fopingensis* Chou et Ma
Aedeagal shaft short, slightly curved, nearly straight
..... *S. nigropunctata* Mahmood
6. Apex of aedeagal preatrium process with furrows
..... *S. arifi* Ghauri
Apex of aedeagal preatrium process without furrows
..... *S. bannaensis* sp. nov.
7. Both side of apex of aedeagal shaft serrated 8

The project was supported by the Key Technologies R&D Program of Guiyang City (No. [2009] 3-0243) and Guizhou Provincial Natural Science Foundation of China (No. [2010] 2063).

* Corresponding author


Received 9 Feb. 2010; accepted 26 Aug. 2010.

- Both side of apex of aedeagal shaft smooth..... 10
 8 Apex of aedeagal shaft truncated *S. shivae* **Dworakowska**
 Apex of aedeagal shaft arced slightly 9
 9 Apex of anal tube process expanded, truncated
 *S. indica* **Ramakrishnan et Menon**
 Apex of anal tube process not expanded, rounded.....
 *S. kamatana* **Viraktamath et Dworakowska**
 10 Anal tube appendage long and fine bent ventrally and tapering to
 apex *S. shishana* **Matsumura**
 Anal tube appendage short and robust, rounded at apex..... 11
 11 Aedeagus preatrial process with base broad and narrowing abruptly
 near apex..... *S. victorena* **Chiang et Knight**


- Aedeagus preatrial process tapering towards to apex.....
 *S. viridis* **Dworakowska**

Singapore bannaensis **sp. nov.** (Figs 1– 8, 18)

Description. Body length: ♂4 mm. Vertex milky yellow, with a black median apical spot and with two small dim spots at sides of coronal suture. Eyes black (Figs 1, 18). Most part of face black, anteclypeus broad, inflated, with a few small microsetae (Fig 2). Pronotum (Figs 1, 18) arcuately rounded at anterior margin, concave at base, fore region whitish, middle



Figs 1– 8 *Singapore bannaensis* **sp. nov.** 1 Head and thorax, dorsal view. 2 Face. 3 Abdominal apodemes. 4 Pygofer, lateral view. 5 Style. 6 Aedeagus, ventral view. 7 Aedeagus, lateral view. 8 Connective.



Figs 9– 17. *Singapon arfi* Ghauri 1985 9. Head and thorax, dorsal view. 10. Face 11. Forewing
12. Abdominal apodemes 13. Pygofer, lateral view. 14. Style 15. Aedeagus, ventral view. 16. Aedeagus
lateral view. 17. Connective

and hind regions brownish. Scutellum transverse impression prominent, curved slightly. Forewing light brown, brochsome field dark brown.

Abdominal apodemes (Fig 3) long, exceeding 7th sternite.

Male genitalia. Anal tube appendage tapering, its baso-ventral margin connected with dorsal apodeme of aedeagus. Pygofer lobe (Fig 4) with numerous microsetae near caudal margin. Subgenital plate (Fig 4) with six long macrosetae at basal angle; four large macrosetae on outer margin medially and numerous

microsetae at apex and outer surface. Style (Fig 5) long and slim, with foot-like apex; preapical lobe poorly developed. Aedeagus (Figs 6– 7) with a long preatrial process, about half apical length of preatrial process very thin; dorsal apodeme dilated distinctly. Aedeagal shaft broad, with pair of long apical processes and longer than length of aedeagal shaft; gonopore apical, ventrad. Connective (Fig 8) almost triangular; two lateral arms long and strong, stem comparatively small and short.

Holotype ♂, China, Yunnan Province.




Fig 18 *Singapore bamaensis* sp. nov. (dorsal view). Fig 19. *Singapore arifi* Ghauri 1985 (dorsal view).

Xishuangbanna (22° 06' N, 100° 55' E), 20 July 2008, coll LI Jian-Da Paratype 1 ♂, same data as holotype

Remarks The new species is similar to *S. arifi* Ghauri (1985) in male genital apparatus but can be distinguished from the latter by its very long lamellate-like apical processes of aedeagal shaft extending beyond the base of aedeagal shaft the aedeagus preatrial process very long expanded at base and quite slender near apex and the face with most region entirely dark (Figs 6–7).

Etymology. The new species is named after its type locality “Banna”.

***Singapore arifi* Ghauri 1985 New record** (Figs 9–17, 19)

Singapore arifi Ghauri 1985 *Türk. Bit. Kor. Derg.* 9: 5, 11.

Description Body length ♂ 4.1–4.3 mm, ♀ 4.5 mm. Vertex (Figs 9, 19) dirty yellow or light brown with a large black apical spot medially, two small dark spots near apical spot not very distinct. Eyes black. Face (Fig 10) frontoclypeus broad, pale and anteclypeus inflated nearly pentagon with numerous small microsetae, just genae and bra black. Pronotum (Figs 9, 19) broad, only anterior margin pale, other areas dark, blackish brown. Scutellum (Fig 9) also dark, with transverse impression curved slightly, arc like, nearer to apex than to base. Forewing (Fig 11) brownish, brochosme field dark.

Abdominal apodemes (Fig 12) long extending to hind margin of 6th sternite.

Male genitalia Anal tube appendage well developed, connecting with dorsal apodeme of aedeagus. Pygofer lobe (Fig 13) with few microsetae around distal margin and lots of microtrichia on lateral surface. Subgenital plate (Fig 13) gently curved dorsad in lateral view, extending beyond caudal margin of pygofer, with several macrosetae at base and five long macrosetae on outer-lateral surface. Style (Fig 14) long and slender, its apex elongate, foot-like near apex with transverse striations. Aedeagus (Figs 15–16) well developed, with a long preatrial process almost reaching beyond apex of shaft, about half apical length of preatrial process with distinct striations and sculptures along both lateral margins. Dorsal apodeme slightly expanded in lateral view and bifurcate apically, preatrium slender. Connective (Fig 17) about triangular, two lateral arms broad and strong stem robust with a longitudinal internal ridge.

Material examined 7 ♂♂, 1 ♀, China Yunnan Prov. Yingjiang Tongbiguan (24° 41' N, 97° 56' E; alt 1 400–1 500 m), 20 July 2002, coll LI Zi Zhong.

Distribution India, China (Yunnan).

Host plant *Derris robusta*.

Acknowledgements We thank Dr Dmitriy D. A. Section for Biodiversity, Illinois Natural History Survey, USA, for providing some literatures.

REFERENCES

Chiang C. C. and Knight W. J. 1990 Studies on taiwanese Typhlocybinae (Homoptera: Cicadellidae) (IV) tribe Erythroneurini *Bulletin of the National Museum of Natural Science*, 2: 240–244

Chou I and Ma N. 1981 On some new species and new records of Typhlocybinae from China (Homoptera: Cicadellidae). *Entomotaxonomica*, 3 (3): 196–197 [周尧, 马宁, 1981. 中国小叶蝉亚科的新种和新纪录 (同翅目: 叶蝉科). 昆虫分类学报, 3 (3): 196–197]

Dworakowska I. 1970 On the genera *Asimilia* Zaidv. and *Singapora* Mahm. with the description of two new genera (Auchenorrhyncha: Cicadellidae: Typhlocybinae). *Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques*, 18 (12): 759–760, 762, 764.

Dworakowska I. and Sohi A. S. 1978 On some Typhlocybinae with remarks on synonymy of Indian species (Auchenorrhyncha: Cicadellidae). *Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques*, 26 (1): 39

Dworakowska I. 1983 The leafhopper *Ranbara* gen. nov. (Cicadellidae: Typhlocybinae) and remarks on *Singapora* Mahm. with description of one new species from Thailand. *Thai Journal of Agricultural Science*, 16: 118–123.

Ghauri M. S. K. 1985. A new species of *Singapora* Mahmood attacking *Derris robusta* in Assam, with a key to the species (Homoptera: Cicadelloidea). *Turkiye Bitki Korunma Dergisi*, 9: 3–11.

Mahmood S. H. 1967. A study of the typhlocybinae genera of the Oriental Region (Thailand, the Philippines and adjoining areas). *Pacific Insects Monograph*, 12: 19–21.

新小叶蝉属一新种及一新纪录种记述（半翅目，叶蝉科，小叶蝉亚科）

宋月华^{1, 2} 李子忠^{1*}

- 1 贵州山地农业病虫害重点实验室，贵州大学昆虫研究所 贵阳 550025
- 2 贵州师范大学中国南方喀斯特研究院 贵阳 550001

摘 要 记述采自中国云南省新小叶蝉属 1 新种，版纳新小叶蝉 *Singapora bannaensis* sp. nov. 及 1 中国新纪录种，阿氏新小叶蝉 *Singapora arfi* Ghauri, 1985。新种区别于属内其它种的特征在于：阳茎端部的刺突很长，其末端已超过阳茎干基部水平；阳茎前腔突基部膨胀，近中部骤细；腹内突长，伸达

第 7 腹节。
词源：新种种名取自采集地地名“版纳”。
同时给出该属世界种类检索表，模式标本保存在贵州大学昆虫研究所。

关键词 半翅目，叶蝉科，小叶蝉亚科，新小叶蝉属，新种，中国。
中图分类号 Q969.35

* 通讯作者。